

ELSEVIER

Carbohydrate Research 269 (1995) C9-C10

Author index

Abbott, C. 1 Abe, Y. 43 An, J. 303

Baczko, K. 79 Baggio, R.F. 99 Banoub, J. 79 Bauer, C.J. 139 Berić, O. 343 Besemer, A.C. 89 Brade, H. C1 Brouillette, J.N. 201

Campos-Portuguez, S. 157 Carlson, R.W. 303 Cases, M.R. 333 Cerezo, A.S. 333 Chai, W. 111, 139 Chida, N. 53 Choy, Y.M. 295 Colquhoun, I.J. 319 Csanádi, J. 343

Defernez, M. 319 De Lederkremer, R.M. 99 De Nooy, A.E.J. 89 Di Nardo, C. 99 Dutton, G.G.S. 295

El Ashry, E.S.H. 349 El Habrouk, M. 349 Elvebak II, L.E. 1 Endo, T. 273, 369

Falshaw, R. 183 Fronczek, F.R. 191 Fujiwara, M. 43 Furneaux, R.H. 183 Garland, M.T. 99 Glushka, J. 303 Gray, G.R. 1, 167

Haasnoot, C.A.G. 17 Harata, K. 43 Hashimoto, H. 273 Helander, A. 125 Herbat, R. 29 Hicks, K.B. 201 Holst, O. C1 Hounsell, E.F. 139

Ijima, H. 227 Irwin, P.L. 201

Jachymek, W. 125 Jones, C. 175

Kajihara, Y. 273 Kenne, L. 125 Kobayashi, S. 369 Kochanowski, H. 157 Kodama, H. 273

Lankhorst, P.P. 17 Lawson, A.M. 111, 139 Lee, C.K. 167 Lee, Y. 191 Leslie, M.R. 295 Lugowski, C. 125

Macindoe, W.M. 227 Mayer, H. 157 Miljković, D. 343 Morris, V.J. 319 Mousaad, A. 349 Müller-Loennies, S. C1 Nagai, T. 369 Nakahara, Y. 227 Neighbors, S.M. 259 Niedziela, T. 125 Nimmich, W. C1 Nugier-Chauvin, C. 79

Ogawa, S. 53 Ogawa, T. 227 Ohbu, K. 43

Paibir, S.G. 259 Parolis, H. 295 Parolis, L.A.S. 295 Petersson, C. 125 Pfannemüller, B. 29 Plusquellec, D. 79 Popsavin, M. 343 Popsavin, V. 343

Rademann, J. 217 Rashed, N. 349 Rosankiewicz, J.R. 111

Saenger, W. 29 Schmidt, R.R. 217 Shalaby, M.A. 191 Shashkov, A.S. 157 Smeets, J.W.H. 17 Soine, W.H. 259 Steiner, T. 29 Stortz, C.A. 333 Streeter, J.G. 303 Süsskind, M. C1

Taki, T. 53 Tarelli, E. 359 Thibault, P. 79

Ueda, H. 369

Van Bekkum, H. 89 Varela, O. 99 Vega, D.R. 99

Wall, S. 1 Wheeler, S.F. 359

Yokoi, S. 53 Yokota, A. 157 Yoshikawa, M. 53 Younathan, E.S. 191

SUBJECT INDEX

Acetan, 319

Acetobacter xylinum, 319

O-Acetylated cellulose by the reductive-cleavage method, analysis of positions of substitution of O-acetyl groups in partially, 167

Acetylated or benzoylated 1,5-anhydro-Dgalactitol, authentic standards for the reductive-cleavage method. The positional isomers of partially methylated and. 1

O-Acetyl groups in partially O-acetylated cellulose by the reductive-cleavage method, analysis of positions of substitution of, 167

N-Acetylneuraminic acid and 3-deoxy-D-manno-2-octulosonic acid (Kdo), synthesis of carbocyclic analogues, 53

6-O-Acyl and 6,6'-di-O-acylsucroses, a new synthesis, 79

Alditols, separation and quantitation of enantiomeric galactoses and their mono-O-methyl ethers as their diasteromeric acetylated 1deoxy-1-(2-hydroxypropylamino), 333

Aldosulose bis(phenylhydrazones), isopropylidenation of, 349

Algal galactans by methylation and reductive partial-hydrolysis, the structural analysis of disaccharides from red, 183

Aliphatic chain packing, 29 Alkyl-gluconamides, 29 Amphiphilic molecules, 29

Bacterial polysaccharides, 319

Barbital, phenobarbital, metharbital, and mephobarbital, synthesis of N-β-D-glucopyranosyluronate derivative of, 259

Benzoylated 1,5-anhydro-p-galactitol, authentic standards for the reductive-cleavage method. The positional isomers of partially methylated and acetylated or, 1

Biantennary penta- and hepta-saccharides having two 6-deoxy-p-galactose residues at the nonreducing end and evaluation of 6-deoxyD-galactosyl transfer to glycoprotein using bovine β -(1 \rightarrow 4)-galactosyltransferase and UDP-6-deoxy-D-galactose, enzymic transfer of 6-modified residues: Synthesis of, 273

Bilaver, 43

Blood group A type glycopeptide present in human blood mucin, stereoselective total synthesis of a, 227

Bradyrhizobium species within soybean nodules, the structure of the novel polysaccharide produced by, 303

Bromination of sugar enones and enonolactones, 99

Capsular polysaccharide, 295

Capsular polysaccharide from Streptococcus pneumoniae serotype 10A, full assignment of the NMR spectrum, 175

Carbocyclic analogues of 3-deoxy-p-manno-octulosonic acid (Kdo) and N-acetylneuraminic acid, synthesis, 53

Cellulose by the reductive-cleavage method, analysis of positions of substitution of Oacetyl groups in partially O-acetylated, 167

Chlorogenic acid: water activity dependence, cyclomaltoheptaose (β-cyclodextrin) and hydroxyethyl-substituted β-cyclodextrin inclusion complex formation with, 201

Conformational analysis, 17

Core/lipid A region of the lipopolysaccharide from Klebsiella pneumoniae rough mutant R20/O structural investigation, C1

Crystal packing, 29 Crystal structure, 43

Cyanogenic glycoside, 17

(η-Cyclodextrin), isolation, purification and characterization of cyclomaltododecaose, 369

Cyclomaltododecaose (η-cyclodextrin), isolation, purification and characterization of, 369

Cyclomaltoheptaose (β-cyclodextrin) and hydroxyethyl-substituted β-cyclodextrin inclu-

- sion complex formation with chlorogenic acid: water activity dependence, 201
- 6-Deoxy-D-galactose residues at the nonreducing end and evaluation of 6-deoxy-D-galactosyl transfer to glycoprotein using bovine β-(1 → 4)-galactosyltransferase and UDP-6-deoxy-D-galactose, enzymic transfer of 6-modified residues: Synthesis of biantennary penta- and hepta-saccharides having two, 273
- 6-Deoxy-D-galactosyl transfer to glycoprotein using bovine β-(1 → 4)-galactosyltransferase and UDP-6-deoxy-D-galactose, enzymic transfer of 6-modified residues: Synthesis of biantennary penta- and hepta-saccharides having two 6-deoxy-D-galactose residues at the nonreducing end and evaluation of, 273
- 3-Deoxy-D-manno-2-octulosonic acid (Kdo) and N-acetylneuraminic acid, synthesis of carbocyclic analogues, 53
- Diaminopyranoses forming the repeating unit, the structure of the O-specific polysaccharide from *Thiobacillus* sp. IFO 14570, with three different, 157
- Diasteromeric acetylated 1-deoxy-1-(2-hydroxypropylamino)alditols, separation and quantitation of enantiomeric galactoses and their mono-O-methyl ethers as their, 333
- Disaccharides from red algal galactans by methylation and reductive partial-hydrolysis, the structural analysis of, 183
- 3D structure, 17
- Enones and enonolactones, bromination of sugar, 99
- Enonolactones, bromination of sugar enones, 99 Escherichia coli, 295
- Galactans by methylation and reductive partialhydrolysis, the structural analysis of disaccharides from red algal, 183
- D-Galactitol, authentic standards for the reductive-cleavage method. The positional isomers of partially methylated and acetylated or benzoylated 1,5-anhydro-, 1
- Galactoses and their mono-O-methyl ethers as their diasteromeric acetylated 1-deoxy-1-(2hydroxypropylamino)alditols, separation and quantitation of enantiomeric, 333
- β-(1→4)-Galactosyltransferase and UDP-6-deoxy-D-galactose, enzymic transfer of 6-modified residues: Synthesis of biantennary penta- and hepta-saccharides having two 6deoxy-D-galactose residues at the nonreducing end and evaluation of 6-deoxy-D-

- galactosyl transfer to glycoprotein using bovine, 273
- N-β-D-Glucopyranosyluronate derivatives of barbital, phenobarbital, metharbital, and mephobarbital, synthesis of, 259
- Glucosidase, 17
- D-Glucose, an alternative synthesis of (+)-epiallo-muscarine from, 343
- Glycopeptide present in human blood mucin, stereoselective total synthesis of a blood group A type, 227
- Glycoprotein using bovine β-(1 → 4)-galactosyltransferase and UDP-6-deoxy-D-galactose, enzymic transfer of 6-modified residues: Synthesis of biantennary penta- and heptasaccharides having two 6-deoxy-D-galactose residues at the nonreducing end and evaluation of 6-deoxy-D-galactosyl transfer to, 273
- Glycosaminoglycan, 111
- Glycosurfactant, 43
- Glycosylated hexapeptide of human sialophorin, solid-phase synthesis using the trichloroacetimidate method, 217
- Hafnia alvei strain 1192 lipopolysaccharide, structural studies of the O-specific chain and a core hexasaccharide, 125
- Heparin, characterisation by LSI-MS and ¹H NMR spectroscopy of oligosaccharides of porcine intestinal, 139
- Heterodendrin, 17
- ¹H NMR spectroscopy and liquid secondary ion mass spectrometry, characterisation of oligosaccharides of porcine intestinal heparin, 139
- Human blood mucin, stereoselective total synthesis of a blood group A type glycopeptide present in, 227
- Hydrogen bond, 43
- Hydrogen bonding, 29
- Inclusion complex formation with chlorogenic acid: water activity dependence, cyclomaltoheptaose (β -cyclodextrin) and hydroxyethylsubstituted β -cyclodextrin, 201
- Interdigitated structure, 43
- Isopropylidenation of aldosulose bis(phenylhydrazones), 349
- K43 antigen, 295
- Klebsiella pneumoniae rough mutant R20/O1⁻, structural investigation on the carbohydrate backbone of the lipopolysaccharide, C1
- Lipopolysaccharide from Klebsiella pneumoniae rough mutant R20/O1⁻, structural investigation of the carbohydrate backbone, C1

- Lipopolysaccharide, structural studies of the Ospecific chain and a core hexasaccharide of Hafnia alvei strain 1192, 125
- Liquid secondary ion mass spectrometry (LSIMS), 111
- D-Mannitol. Comparison of NMR spectral results for the solid state and solution with those of the X-ray structural determination, synthesis of 3,4-di-O-acetyl-2,5-anhydro-1,6-dideoxy-1,6-diiodo-, 191
- Mephobarbital, synthesis of N-β-D-glucopyranosyluronate derivatives of barbital, phenobarbital, metharbital, and, 259
- Metharbital, and mephobarbital, synthesis of N-β-D-glucopyranosyluronate derivatives of barbital, phenobarbital, 259
- Methylated and acetylated or benzoylated 1,5anhydro-p-galactitol, authentic standards for the reductive-cleavage method. The positional isomers of partially, 1
- Methylation and reductive partial-hydrolysis, the structural analysis of disaccharides from red algal galactans by, 183

MM2, 17

(+)-epiallo-Muscarine, an alternative synthesis from p-glucose, 343

Neoglycolipids, 111

- Nitroxyl radical-mediated oxidation of primary alcohol groups in water-soluble glucans, 89
- NMR spectral results for the solid state and solution with those of the X-ray structural determination, synthesis of 3,4-di-O-acetyl-2,5-anhydro-1,6-dideoxy-1,6-diiodo-D-mannitol. Comparison of, 191

NMR spectroscopy, 295, 319 Nonionic surfactant, 43

Oligosaccharides of porcine intestinal heparin, characterisation by LSI-MS and ¹H NMR spectroscopy, 139

Oxymercuration, 111

- Phenobarbital, metharbital, and mephobarbital, synthesis of N-β-p-glucopyranosyluronate derivatives of barbital, 259
- Polysaccharide from Streptococcus pneumoniae serotype 10A, full assignment of the NMR spectrum of the capsular, 175
- Polysaccharide produced by *Bradyrhizobium* species within soybean nodules, the structure of the novel, 303
- Primary alcohol groups oxidation in water soluble glucans with nitroxyl radicals, 89

- Reductive-cleavage method, analysis of positions of substitution of O-acetyl groups in partially O-acetylated cellulose by the, 167
- Reductive-cleavage method. The positional isomers of partially methylated and acetylated or benzoylated 1,5-anhydro-p-galactitol, authentic standards for the, 1
- Reductive partial-hydrolysis, the structural analysis of disaccharides from red algal galactans by methylation and, 183
- Solid-phase synthesis of a glycosylated hexapeptide of human sialophorin, using the trichloroacetimidate method, 217
- O-Specific polysaccharide and a core hexasaccharide of the *Hafnia alvei* strain 1192 lipopolysaccharide, structural studies, 125
- O-Specific polysaccharide from Thiobacillus sp. 1FO 14570, with three different diaminopyranoses forming the repeating unit, the structure of the, 157
- Stereospecific synthesis of (+)-epiallo-muscarine from D-glucose, 343
- Streptococcus pneumoniae serotype 10A, full assignment of the NMR spectrum of the capsular polysaccharide, 175
- Structural analysis of disaccharides from red algal galactans by methylation and reductive partial-hydrolysis, the, 183
- Structure of the novel polysaccharide produced by *Bradyrhizobium* species within soybean nodules, the, 303
- Structure of the O-specific polysaccharide from Thiobacillus sp. IFO 14570, with three different diaminopyranoses forming the repeating unit, the, 157
- Substitution of O-acetyl groups in partially Oacetylated cellulose by the reductive-cleavage method, analysis of positions of, 167
- Sucrose and sodium phosphate, the preparation of sucrose monophosphates from dried mixtures of, 359
- Sucrose monophosphates from dried mixtures of sucrose and sodium phosphate, the preparation of, 359

Surfactant, 43

- Synthesis of a blood group A type glycopeptide present in human blood mucin, stereoselective total, 227
- Synthesis of 3,4-di-O-acetyl-2,5-anhydro-1,6-dideoxy-1,6-diiodo-n-mannitol. Comparison of NMR spectral results for the solid state and solution with those of the X-ray structural determination, 191

TLC-LSIMS, 111

Trichloroacetimidate method, use in the solidphase synthesis of a glycosylated hexapeptide of human sialophorin, 217

UDP-6-deoxy-D-galactose, enzymic transfer of 6-modified residues: Synthesis of biantennary penta- and hepta-saccharides having two 6-deoxy-D-galactose residues at the nonreducing end an evaluation of 6-deoxy-D-galactosyl transfer to glycoprotein using bovine β -(1 \rightarrow 4)-galactosyltransferase and, 273

Water-soluble glucans oxidation of primary alcohol groups with nitroxyl radicals, 89

X-ray analysis, 43

X-ray structural determination, synthesis of 3,4-di-O-acetyl-2,5-anhydro-1,6-dideoxy-1,6-diiodo-D-mannitol. Comparison of NMR spectral results for the solid state and solution with those of the, 191

